

## End of Result Set

Generate Collection

L6: Entry 1 of 1

File: USPT

Aug 31, 1999

US-PAT-NO: 5945275

DOCUMENT-IDENTIFIER: US 5945275 A

TITLE: Nematode-extracted anti-inflam protein

DATE-ISSUED: August 31, 1999

## INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Vlasuk; George Phillip	Carlsbad	CA	N/A	N/A
Stanssens; Patrick Eric Hugo	St-Martens-Latem	N/A	N/A	BEX
Messens; Joris Hilda Lieven	Antwerpen	N/A	N/A	BEX
Lauwerijs; Marc Jozef	Haaltert	N/A	N/A	BEX
Laroche; Yves Rene	Brussels	N/A	N/A	BEX
Jaspers; Laurent Stephane	Tervuren	N/A	N/A	BEX
Gansemans; Yannick Georges Jozef	Bredene	N/A	N/A	BEX

US-CL-CURRENT: 435/5; 435/6; 536/23.5

## CLAIMS:

We claim:

1. A recombinant cDNA molecule having a nucleic acid sequence selected from the nucleic acid sequences depicted in FIG. 7A [SEQ. ID. NO. 38], FIG. 7B [SEQ. ID. NO. 39], FIG. 7C [SEQ. ID. NO. 40], FIG. 7D [SEQ. ID. NO. 41], FIG. 7E [SEQ. ID. NO. 42], and FIG. 7F [SEQ. ID. NO. 43].
2. A recombinant cDNA molecule having the nucleic acid sequence depicted in FIG. 9 [SEQ. ID. NO. 47].
3. A recombinant cDNA molecule having a nucleic acid sequence selected from the nucleic acid sequences depicted in FIG. 1 [SEQ. ID. NO. 32] and FIG. 3 [SEQ. ID. NO. 34].
4. A recombinant cDNA molecule characterized as having the nucleic acid sequence depicted in FIG. 9 [SEQ. ID. NO. 47] and made by a method of making a recombinant cDNA encoding a protein, comprising the steps of:
  - (a) isolating a cDNA library from a species of nematode selected from the group consisting of Ancylostoma caninum, Ancylostoma ceylanicum, Ancylostoma duodenale, Necator americanus and Heligmosomoides polygyrus;
  - (b) ligating said cDNA library into a phagemid vector selected from the group consisting of pDONG61 [SEQ. ID. NO. 44], pDONG62 [SEQ. ID. NO. 45] and pDONG63 [SEQ. ID. NO. 46];
  - (c) transforming host cells which are E. coli, strain TG1 with said vector containing said cDNA library;
  - (d) culturing said host cells;
  - (e) infecting said host cells with a helper phage;
  - (f) separating phage containing said cDNA library from said host cells;
  - (g) combining a solution of said phage containing said cDNA library with a solution of biotinylated human factor Xa;
  - (h) contacting streptavidin-coated solid phase with said solution containing said phage containing said cDNA library, and said biotinylated human factor Xa;
  - (i) isolating phage which bind to said streptavidin-coated solid phase; and
  - (j) isolating the recombinant cDNA molecule from phage which bind to said streptavidin-coated solid phase.